antibodies .-online.com

Datasheet for ABIN1389847 anti-Cadherin 4 antibody (AA 151-250) (Biotin)



Overview

Quantity:	100 μL
Target:	Cadherin 4 (CDH4)
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cadherin 4 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human R Cadherin
Isotype:	lgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	Cadherin 4 (CDH4)
Alternative Name:	R Cadherin (CDH4 Products)
Background:	Synonyms: CADH4_HUMAN, Cadherin 4, Cadherin4, Cadherin-4, CDH 4, CDH-4, R CAD, R-CAD,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1389847 | 03/07/2024 | Copyright antibodies-online. All rights reserved. R-cadherin, Retinal Cadherin.

Background: The cadherins are a family of Ca++-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including catenin b, to regulate cadherin function. R-cadherin (for retinal-cadherin, also designated cadherin-4), which was first identified in the retina of chicken, has been shown to be involved in the development of striated muscle and potentially epithelia in addition to its involvement in retinal development.

Pathways:

Cell-Cell Junction Organization, Regulation of Cell Size

Application Details

Restrictions:	For Research Use only
	IHC-F 1:100-500
	IHC-P 1:200-400
Application Notes:	WB 1:300-5000

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months