

[Go to Product page](#)

Datasheet for ABIN1389965

**anti-MTMR14 antibody (AA 161-260) (Alexa Fluor 555)**

## Overview

Quantity:	100 µL
Target:	MTMR14
Binding Specificity:	AA 161-260
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTMR14 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MTMR14
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	MTMR14
Alternative Name:	MTMR14 ( <a href="#">MTMR14 Products</a> )

## Target Details

Background:	<p>Synonyms: C3orf29, Egg derived tyrosine phosphatase homolog, FLJ11546, FLJ22405, FLJ46453, FLJ90311, HCV NS5A transactivated protein 4 splice variant A binding protein 1, HCV NS5A-transactivated protein 4 splice variant A-binding protein 1, hJumpy, jumpy, MTMR 14, MTMR14, MTMR-14, MTMRE_HUMAN, Myotubularin related protein 14, Myotubularin-related protein 14, NS5ATP4ABP1.</p> <p>Background: Myotubularin-related protein 14 (MTMR14), also known as Jumpy, is a myotubularin-related phosphoinositol-3-phosphate (PI3P) phosphatase (1). Mutations in the MTMR14 gene have been associated with centronuclear myopathy (1). MTMR14 deficiency in mice leads to altered calcium homeostasis and muscle disorders (2). MTMR14 has also been shown to play a role in autophagy, a process that is highly regulated by phosphatidylinositides through the type III PI3K, Vps34 (3). MTMR14 was localized to autophagic isolation membranes and early autophagosomes (3). In these studies, MTMR14 inhibited autophagy and mutations of MTMR14 associated with centronuclear myopathy were also defective in autophagy inhibition. In zebrafish, MTMR14 knockdown was shown to increase the number of autophagosomes, suggesting that its activity is associated with an inhibition of autophagy (4).</p>
-------------	--

Pathways:	<a href="#">Inositol Metabolic Process</a>
-----------	--

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months