
# antibodies -online.com





## anti-SH3TC2 antibody (AA 851-950) (Alexa Fluor 555)



Go to Product page

$\sim$					
( )	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	SH3TC2
Binding Specificity:	AA 851-950
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH3TC2 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SH3TC2	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Cow, Pig, Horse, Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	SH3TC2	
Alternative Name:	SH3TC2 (SH3TC2 Products)	
Background:	ckground: Synonyms: KIAA1985, CMT4C, D430044G18Rik, FLJ13605, MNMN, PP12494, RGD13090	

S3TC2\_HUMAN, SH3 domain and tetratricopeptide repeats-containing protein 2, SH3TC2. Background: SH3TC2 (SH3 domain and tetratricopeptide repeats 2) is a 1,288 amino acid protein that contains one SH3 domain and eight TPR repeats. The SH3TC2 gene encodes a protein expressed in Schwann cells of peripheral nerves, and localized to the plasma membrane and to the perinuclear endocytic recycling compartment, suggesting a possible function in myelination and/or in regions of axoglial interactions. The SH3TC2 protein is expressed in adult heart, testis, spinal cord, and brain as well as in fetal brain and liver. Mild mononeuropathy of the median nerve (MNMN) is caused by heterozygous mutation in the SH3TC2 gene. Also, Charcot-Marie-Tooth disease type 4C (CMT4C) is a more severe neuropathy caused by homozygous or compound heterozygous mutation in the SH3TC2 gene. Existing as four alternatively spliced isoforms and containing 18 exons, the SH3TC2 gene is conserved in chimpanzee, dog, cow, mouse, rat, chicken and zebrafish, and maps to human chromosome 5q32.

#### **Application Details**

Application Notes:

Storage Comment:

**Expiry Date:** 

	IF(IHC-F) 1:50-200 IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	

IF(IHC-P) 1:50-200

12 months

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.