



[Go to Product page](#)

Datasheet for ABIN1390086

anti-TCF12 antibody (AA 51-150) (Alexa Fluor 647)

Overview

Quantity:	100 µL
Target:	TCF12
Binding Specificity:	AA 51-150
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TCF12 antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TCF12
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse, Sheep, Horse, Rabbit, Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	TCF12
Alternative Name:	TCF12/HEB (TCF12 Products)

Target Details

Background: Synonyms: bHLHb20, Class B basic helix-loop-helix protein 20, DNA binding protein HTF4, DNA-binding protein HTF4, E box binding protein, E-box-binding protein, HEB, HTF4, HTF4_HUMAN, TCF-12, Tcf12, Transcription factor 12, Transcription factor HTF-4, Transcription factor HTF4.

Background: Differentiation of myogenic cells is regulated by multiple positively and negatively acting factors. One well characterized family of helix-loop-helix (HLH) proteins known to play an important role in the regulation of muscle cell development includes Myo D, myogenin, Myf-5 and Myf-6 (also designated MRF-4 or herculin). Myo D transcription factors form heterodimers with products of a more widely expressed family of bHLH genes, the E family, which consists of at least three distinct genes: E2A, IF2 and HEB. Myo D-E heterodimers bind avidly to consensus (CANNTG) E box target sites that are functionally important elements in the upstream regulatory sequences of many muscle-specific terminal differentiation genes. Both homo- and hetero-oligomers of these proteins are able to distinguish very closely related E box proteins and are believed to play important roles in lineage specific gene expression.

Gene ID: 6938

UniProt: [Q99081](#)

Pathways: [SARS-CoV-2 Protein Interactome](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months