
# antibodies -online.com





## anti-CREB3L3 antibody (AA 201-300) (Biotin)



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL
Target:	CREB3L3
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CREB3L3 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CREB-H
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

## **Target Details**

Target:	CREB3L3
Alternative Name:	CREB-H/CREB3L3 (CREB3L3 Products)

#### Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		$\sim$	<i>.</i>	ч.

Synonyms: cAMP responsive element binding protein 3 like 3, cAMP responsive element-binding protein 3 like protein 3, CREB H, CREB/ATF family transcription factor, CREBH, Cyclic AMP responsive element binding protein 3 like protein 3, cAMP-responsive element-binding protein 3-like protein 3, HYST1481, MGC126553, MGC126557, Transcription factor CREB H, CR3L3\_HUMAN.

Background: CREB3L3 is a 461 amino acid single-pass type II membrane protein that localizes to the endoplasmic reticulum (ER) and, in response to ER stress, is cleaved and translocated to the nucleus. Expressed exclusively in liver, CREB3L3 functions as a transcription factor that, during ER stress, is thought to activate genes that are involved in both the unfolded protein response and the acute phase response (APR). Additionally, CREB3L3 is underexpressed in hepatocellular carcinoma, suggesting a possible role as a tumor suppressor. CREB3L3 functions as a dimer and contains one leucine zipper domain, a KDEL-like sequence and a bZIP domain, through which it conveys its DNA binding ability. Three isoforms of CREB3L3 exist due to alternative splicing events.

Pathways:

Thyroid Hormone Synthesis

#### **Application Details**

$\Delta$ nn	lication	Notae.

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months