

Datasheet for ABIN1390189

**anti-NSF antibody (AA 151-250) (Biotin)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	NSF
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NSF antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NSF
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit,Zebrafish
Purification:	Purified by Protein A.

## Target Details

Target:	NSF
Alternative Name:	NSF ( <a href="#">NSF Products</a> )

## Target Details

Background:	<p>Synonyms: N ethylmaleimide sensitive factor, N ethylmaleimide sensitive factor, N ethylmaleimide sensitive factor like protein, N ethylmaleimide sensitive fusion protein, N ethylmaleimide sensitive fusion protein, N-ethylmaleimide-sensitive fusion protein, NEM sensitive fusion protein, NEM sensitive fusion protein, NEM-sensitive fusion protein, NSF, NSF_HUMAN, SKD 2, SKD2, SKD2, Vesicle fusing ATPase, Vesicle fusing ATPase, Vesicle-fusing ATPase, Vesicular fusion protein NSF, Vesicular fusion protein NSF, Vesicular-fusion protein NSF.</p> <p>Background: Several protein-protein interactions are essential to membrane fusion during endocytosis. Membrane fusion requires interaction among SNARE1 proteins associated with both donor and acceptor membranes (1,2). Following membrane fusion, the -SNAP cytoplasmic adapter protein binds to the SNARE complex. N-ethylmaleimide-sensitive factor (NSF), a hexameric ATPase, then associates with the -SNAP/SNARE complex to mediate SNARE disassembly during membrane fusion (3,4). The ATPase activity of NSF induces a conformational change in the -SNAP/SNARE complex that leads to its dissociation from the membrane, membrane fusion and eventual recycling of the SNARE complex for subsequent membrane fusion (3,4).</p>
-------------	--

## Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:                      Store at -20°C for 12 months.

---

Expiry Date:                              12 months