## antibodies .- online.com





Datasheet for ABIN1390394

## anti-HAS2 antibody (FITC)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

Background:

| Quantity:         | 100 μL  |  |
|-------------------|---|--|
| Target:           | HAS2  |  |
| Reactivity:       | Human, Mouse, Rat   |  |
| Host:             | Rabbit  |  |
| Clonality:        | Polyclonal  |  |
| Conjugate:        | This HAS2 antibody is conjugated to FITC  |  |
| Application:      | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |
| Product Details   |   |  |
| Immunogen:        | KLH conjugated synthetic peptide derived from human HAS2/Hyaluronan synthase 2  |  |
| Isotype:          | IgG   |  |
| Cross-Reactivity: | Human, Mouse, Rat   |  |
| Purification:     | Purified by Protein A.  |  |
| Target Details    |   |  |
| Target:           | HAS2  |  |
| Alternative Name: | Has2 (HAS2 Products)  |  |
|                   |   |  |

2, Hyaluronic acid synthase 2.

Synonyms: HA synthase 2, has2, HAS2\_HUMAN, Hyaluronan synthase 2, Hyaluronate synthase

Background: HAS1, HAS2 and HAS3 are HA Synthase proteins that synthesize HA (Hyaluronan

or hyaluronic acid). The extracellular matrix in most vertebrates express HA, which is a high

molecular weight linear polysaccharide composed of alternating glucuronic acid and N-acetylglucosamine residues linked by i²-1,3 and i²-1,4 glycosidic bonds. The three HAS genes show distinct patterns of expression during development and their protein products play significantly different roles in the formation of the HA matrix. Both HAS1 and HAS2 synthesise high molecular-weight HA, whereas HAS3 produces lower molecular weight HA. The expression of the three HAS isoforms is more prominent in growing cells than in resting cells and is differentially regulated by various stimuli suggesting distinct functional roles of the three proteins. HAS2 mRNA shows predominant expression in chondrocytes and cartilage. The human HAS2 gene maps to chromosome 8q24.12.

Pathways:

Glycosaminoglycan Metabolic Process

## **Application Details**

Application Notes: IF(IHC-P): (1:50-200)

Optimal working dilution should be determined by the investigator.

Restrictions: Fe

For Research Use only

## Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.             |  |
| Preservative:      | Sodium azide   |  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                      |  |
| Expiry Date:       | 12 months  |  |