



[Go to Product page](#)

Datasheet for ABIN1390411
anti-HOXA4 antibody (AA 151-250) (Biotin)

Overview

Quantity:	100 µL
Target:	HOXA4
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HOXA4 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HOXA4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	HOXA4
Alternative Name:	HOXA4 (HOXA4 Products)

Target Details

Background:	<p>Synonyms: Dfd like protein, Homeo box A4, Homeobox A4, Homeobox protein Hox-1.4, Homeobox protein Hox-1D, Homeobox protein Hox-A4, Hox 1.4 like protein, hox-1.4, hox-1d, HOX1, HOX1D, HOXA4, HXA4_HUMAN.</p> <p>Background: The Hox homeobox genes encode proteins that are transcriptional regulators with an established role in embryonic development. HoxA4 (homeobox A4), also known as HOX1D or HOX1, is a 320 amino acid protein that localizes to the nucleus and contains one homeobox DNA-binding domain. Expressed in the embryonic nervous system, HoxA4 functions as a sequence-specific DNA-binding transcription factor that is part of a regulatory mechanism that provides cells with positional identities during development. Via its ability to bind DNA, HoxA4 plays an important role in the regulation of gene expression, as well as morphogenesis and differentiation. The gene encoding HoxA4 maps to human chromosome 7, which houses over 1,000 genes and comprises nearly 5 % of the human genome. Defects in some of the genes localized to chromosome 7 have been linked to Osteogenesis imperfecta, Williams-Beuren syndrome, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome.</p>
-------------	--

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months