

[Go to Product page](#)

Datasheet for ABIN1390495

**anti-Prostacyclin Receptor antibody (AA 101-200) (Biotin)**

## Overview

Quantity:	100 µL
Target:	Prostacyclin Receptor (PTGIR)
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prostacyclin Receptor antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Prostaglandin I2 Receptor
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	Prostacyclin Receptor (PTGIR)
Alternative Name:	Prostaglandin I2 Receptor ( <a href="#">PTGIR Products</a> )

## Target Details

Background:	<p>Synonyms: IP, PGI receptor, PGI2 receptor, PI2R_HUMAN, PRIPR, Prostacyclin receptor, Prostaglandin I2 receptor, Prostanoid IP receptor, PTGIR.</p> <p>Background: Cyclooxygenases metabolize arachidonate to five primary prostanoids: PGE2, PGF2?, PGI2, TXA2 and PGD2. These lipid mediators interact with specific members of G protein-coupled prostanoid receptors, designated EP, FP, IP, TP and DP, respectively. The IP Receptor binds prostacyclin, PGI2, the main pro-stanoid synthesized by vascular tissues. Upon binding to the IP Receptor, prostacyclin activates adenylate cyclase primarily through the Gas protein. The gene encoding the human IP Receptor is located on chromosome 19. It is expressed as a glycosylated and phosphorylated protein, which is abundantly expressed in vascular tissues such as aorta, lung, atrium and ventricle, as well as in kidney, thymus, spleen and neurons.</p>
Pathways:	<a href="#">cAMP Metabolic Process</a> , <a href="#">Platelet-derived growth Factor Receptor Signaling</a> , <a href="#">Thromboxane A2 Receptor Signaling</a>

## Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months