antibodies -online.com





anti-SNAPIN antibody (AA 51-136) (FITC)



Overview

Quantity:	100 μL	
Target:	SNAPIN	
Binding Specificity:	AA 51-136	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SNAPIN antibody is conjugated to FITC	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SNAP25BP	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	SNAPIN	
Alternative Name:	SNAPIN/SNAP25BP (SNAPIN Products)	
Background:	Synonyms: SNAP associated protein, SNAP-25-binding protein, SNAP-associated protein,	

SNAP25BP, SNAPAP, SNAPIN, SNAPN_HUMAN, SNARE associated protein snapin, SNARE-
associated protein Snapin, Synaptosomal associated protein 25 binding protein, Synaptosomal-
associated protein 25-binding protein, OTTHUMP0000035157.

Background: The protein encoded by this gene is a coiled-coil-forming protein that associates with the SNARE (soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptor) complex of proteins and the BLOC-1 (biogenesis of lysosome-related organelles) complex. Biochemical studies have identified additional binding partners. As part of the SNARE complex, it is required for vesicle docking and fusion and regulates neurotransmitter release. The BLOC-1 complex is required for the biogenesis of specialized organelles such as melanosomes and platelet dense granules. Mutations in gene products that form the BLOC-1 complex have been identified in mouse strains that are models of Hermansky-Pudlak syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012].

Gene ID: 23557

Pathways: Synaptic Membrane, Synaptic Vesicle Exocytosis

Application Details

Application Notes:	IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months