# antibodies .- online.com







## anti-Utrophin antibody (Biotin)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	Utrophin (UTRN)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Utrophin antibody is conjugated to Biotin
Application:	ELISA

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Utrophin
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	Utrophin (UTRN)	
Alternative Name:	Utrophin (UTRN Products)	
Background:	Synonyms: DMDL, DRP 1, DRP, DRP-1, DRP1, Dystrophin like protein, Dystrophin related protein 1, Dystrophin related protein, Dystrophin-related protein 1, FLJ23678, UTRN, UTRO_HUMAN, Utrophin homologous to dystrophin, Utrophin	
	Background: Dystrophin and utrophin are related structural, Actin-binding proteins that are	

involved in anchoring the cytoskeleton to the plasma membrane. Dystrophin is the protein product of the Duchenne/Becker muscular dystrophy gene. Dystrophin expression is found in muscle and brain tissues, where it is localized to the inner surface of the plasma membrane. It has been speculated that alternative splicing of the carboxy terminus allows dystrophin to interact with a variety of proteins. Research has shown that the loss of dystrophin-associated proteins in Duchenne afflicted muscle is due to the absence of dystrophin rather than to muscle degradation and that the lack of dystrophin results in the loss of linkage between the cytoskeleton and the extracellular matrix. Evidence suggests that the upregulation of utrophin can reduce the dystrophic pathology.

Gene ID:

7402

Pathways:

Skeletal Muscle Fiber Development

## **Application Details**

Application Notes:

IHC-P: 1:100-500

Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months