antibodies -online.com





Datasheet for ABIN1391102

anti-MC2R antibody (AA 67-105) (FITC)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	MC2R	
Binding Specificity:	AA 67-105	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MC2R antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	ogen: KLH conjugated synthetic peptide derived from human MC2 receptor	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog	
Purification:	Purified by Protein A.	

Target Details

Target:	MC2R
Alternative Name:	ACTHR (MC2R Products)
Synonyms: ACTH receptor, ACTH-R, ACTHR, ACTHR_HUMAN, Adrenocorticotropic hormone	

receptor, Adrenocorticotropin receptor, MC2-R, MC2R, Melanocortin 2 receptor, Melanocortin	
receptor 2.	

Background: MC2R encodes one member of the five-member G-protein associated melanocortin receptor family. Melanocortins (melanocyte-stimulating hormones and adrenocorticotropic hormone) are peptides derived from pro-opiomelanocortin (POMC). MC2R is selectively activated by adrenocorticotropic hormone, whereas the other four melanocortin receptors recognize a variety of melanocortin ligands. Mutations in MC2R can result in familial glucocorticoid deficiency.

Gene ID: 4158

Pathways: cAMP Metabolic Process

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months