antibodies -online.com





anti-MC4R antibody (AA 1-50) (AbBy Fluor® 488)



Image



Go to Product page

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 μL
Target:	MC4R
Binding Specificity:	AA 1-50
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MC4R antibody is conjugated to AbBy Fluor® 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MC4 Receptor
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	MC4R
Alternative Name:	MC4 Receptor (MC4R Products)
Background:	Synonyms: MC4 R, MC4-R, MC4R, MC4R_HUMAN, Melanocortin 4 receptor, Melanocortin

	receptor 4.	MGC126851.	MGC138197.
--	-------------	------------	------------

Background: The melanocortin family comprises the Alpha-, Beta- and Gamma- melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane G protein-coupled proteins that activate adenylyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.

Gene ID:	4160
UniProt:	P32245


Pathways: Hormone Transport, cAMP Metabolic Process, Feeding Behaviour

Application Details

Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months



Flow Cytometry

Image 1. 293T cells probed with MC4 Receptor Polyclonal Antibody, ALEXA FLUOR® 488 Conjugated (bs-11417R-A488) at 1:100 for 30 minutes compared to control cells (blue) and isotype control (orange).