

Datasheet for ABIN1391274

anti-MED17 antibody (AA 181-280) (AbBy Fluor® 647)



_	verview				
	1//	r	1//	\triangle	۸/
	V		VI		/ V

Background:

Overview			
Quantity:	100 μL		
Target:	MED17		
Binding Specificity:	AA 181-280		
Reactivity:	Human, Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This MED17 antibody is conjugated to AbBy Fluor® 647		
Application:	Western Blotting (WB), Flow Cytometry (FACS)		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human CRSP77		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse		
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit		
Purification:	Purified by Protein A.		
Target Details			
Target:	MED17		
Alternative Name:	CRSP77 (MED17 Products)		
D I	O ODODE ODODET DDIDOO TDADOO MARIE (DMARIE)		

Synonyms: CRSP6, CRSP77, DRIP80, TRAP80, Mediator of RNA polymerase II transcription

subunit 17, Activator-recruited cofactor 77 kDa component, ARC77, Cofactor required for Sp1 transcriptional activation subunit 6, CRSP complex subunit 6, Mediator complex subunit 17, Thyroid hormone receptor-associated protein complex 80 kDa component, Transcriptional coactivator CRSP77, Vitamin D3 receptor-interacting protein complex 80 kDa component, MED17, DRIP77

Background: Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Gene ID: 9440

UniProt: Q9NVC6

Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway, Stem Cell Maintenance, Regulation

of Lipid Metabolism by PPARalpha

Application Details

Application Notes: FCM 1:20-100

Restrictions: For Research Use only

Handling Format: Liquid Concentration: $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. -20 °C Storage: Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. **Expiry Date:** 12 months