antibodies .- online.com





anti-MED17 antibody (AA 181-280) (Biotin)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	MED17
Binding Specificity:	AA 181-280
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED17 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRSP77
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MED17	
Alternative Name:	CRSP77 (MED17 Products)	
Background:	Synonyms: CRSP6, CRSP77, DRIP80, TRAP80, Mediator of RNA polymerase II transcription	

subunit 17, Activator-recruited cofactor 77 kDa component, ARC77, Cofactor required for Sp1 transcriptional activation subunit 6, CRSP complex subunit 6, Mediator complex subunit 17, Thyroid hormone receptor-associated protein complex 80 kDa component, Transcriptional coactivator CRSP77, Vitamin D3 receptor-interacting protein complex 80 kDa component, MED17, DRIP77

Background: Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Gene ID: 9440

UniProt: Q9NVC6

Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway, Stem Cell Maintenance, Regulation

of Lipid Metabolism by PPARalpha

Application Details

Restrictions:

Application Notes: WB 1:300-5000

For Research Use only

Handling Format: Liquid Concentration: $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. -20 °C Storage: Store at -20°C for 12 months. Storage Comment: **Expiry Date:** 12 months