antibodies -online.com





Datasheet for ABIN1391384

anti-DSCAM antibody (AA 301-400) (FITC)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	DSCAM
Binding Specificity:	AA 301-400
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DSCAM antibody is conjugated to FITC
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Dscam	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	DSCAM
Alternative Name:	Dscam (DSCAM Products)

Target Details

•		
Background:	Synonyms: CHD2, CHD2-42, CHD2-52, Down syndrome cell adhesion molecule, DSCAM	
	Background: Cell adhesion molecule that plays a role in neuronal self-avoidance. Promotes	
	repulsion between specific neuronal processes of either the same cell or the same subtype of	
	cells. Mediates within retinal amacrine and ganglion cell subtypes both isoneuronal self-	
	avoidance for creating an orderly dendritic arborization and heteroneuronal self-avoidance to	
	maintain the mosaic spacing between amacrine and ganglion cell bodies. Receptor for netrin	
	required for axon guidance independently of and in collaboration with the receptor DCC. In	
	spinal chord development plays a role in guiding commissural axons projection and pathfinding	
	across the ventral midline to reach the floor plate upon ligand binding. Enhances netrin-induced	
	phosphorylation of PAK1 and FYN. Mediates intracellular signaling by stimulating the activation	
	of MAPK8 and MAP kinase p38.	
Gene ID:	1826	
UniProt:	060469	
Pathways:	Regulation of Cell Size	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months