

[Go to Product page](#)

Datasheet for ABIN1391439

**anti-NAV1 antibody (AA 1426-1475) (Alexa Fluor 350)**

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | NAV1   |
| Binding Specificity: | AA 1426-1475   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This NAV1 antibody is conjugated to Alexa Fluor 350  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Neuron navigator 1 |
| Isotype:              | IgG  |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Pig, Chicken                                   |
| Purification:         | Purified by Protein A.   |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | NAV1   |
| Alternative Name: | Neuron navigator 1 ( <a href="#">NAV1 Products</a> )                                     |
| Background:       | Synonyms: Nav 1, Nav1, POMFIL 3, POMFIL3, Pore membrane and/or filament interacting like |

## Target Details

protein 3, Protein Steerin 1, Protein Steerin1, Steerin 1, Steerin1, unc-53 homolog 1, DKFZp781D0314, FLJ12560, FLJ14203, mNav1, NAV1\_HUMAN.

Background: Neuron navigator 1 is a 1877 amino acid cytoplasmic protein that is involved in neuronal migration. Neuron navigator 1 is widely expressed at low levels, though highest expression is found in both adult and fetal nervous tissue. Through interaction with tubulin, Neuron navigator 1 associates with a subset of microtubule plus ends present in the growth cone. Overexpression of Neuron navigator 1 leads to microtubule bundling, whereas a reduction of its levels causes loss of directionality in the migration of pontine cell leading processes. There are seven isoforms of Neuron navigator 1 that are produced as a result of alternative splicing events.

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months