-online.com antibodies

Datasheet for ABIN1391448 anti-Nogo B Receptor antibody (AA 167-210) (Alexa Fluor 647)



Overview

Quantity:	100 µL	
Target:	Nogo B Receptor (NUS1)	
Binding Specificity:	AA 167-210	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Nogo B Receptor antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NGBR/Nogo B receptor	
lsotype:	lgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	
Target Details		
Target:	Nogo B Receptor (NUS1)	
Alternative Name:	NGBR/Nogo B receptor (NUS1 Products)	
Background:	Synonyms: C6orf68, NGBR, nuclear undecaprenyl pyrophosphate synthase 1 homolog, NUS1,	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1391448 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	NCPD HI IMAN Dobydrodolichyl dinhoanhata gynthaga geranlay gyhynit NUC1. Cia		
	NGBR_HUMAN, Dehydrodolichyl diphosphate synthase complex subunit NUS1, Cis-		
	prenyltransferase subunit NgBR, Nogo-B receptor.		
	Background: Nogo is an oligodendrocyte-specific member of the Reticulon family and is a		
	component of CNS white matter that inhibits axon outgrowth, induces collapse of growth		
	cones of chick dorsal root ganglion cells, and inhibits the spreading of 3T3 fibroblasts. Nogo is		
	expressed by oligodendrocytes but not by Schwann cells and associates primarily with the		
	endoplasmic reticulum. Nogo exists in three different splice forms, Nogo-A, -B and -C. NgBR		
	(Nogo-B receptor), also known as nuclear undecaprenyl pyrophosphate synthase 1 homolog, is		
	a 293 amino acid single-pass type I membrane protein that acts as a specific receptor for the		
	amino-terminus of Nogo-B. Through this interaction, NgBR is involved in the regulation of		
	vascular remodeling and angiogenesis. NgBR also enhances Niemann-Pick type C2 protein		
	(NPC2) stabilization. Knockdown of NgBR mRNA leads to decreased NPC2 levels, which results		
	in the hallmarks of NPC2 mutation: increased intracellular cholesterol accumulation and a loss		
	of sterol sensing.		
Gene ID:	116150		
UniProt:	Q96E22		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Otavaaa			

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1391448 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

-20 °C

Storage:

lond	lina
land	
10110	

Storage Comment:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1391448 | 03/07/2024 | Copyright antibodies-online. All rights reserved.