

Datasheet for ABIN1391503

## **anti-SEMA4B antibody (AA 165-210) (Biotin)**



[Go to Product page](#)

### Overview

Quantity:	100 µL
Target:	SEMA4B
Binding Specificity:	AA 165-210
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SEMA4B antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SEMA4B
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	SEMA4B
Alternative Name:	SEMA4B ( <a href="#">SEMA4B Products</a> )
Background:	Synonyms: Sema domain immunoglobulin domain Ig transmembrane domain TM and short

Target Details

cytoplasmic domain semaphorin 4B, Sema domain immunoglobulin domain Ig transmembrane domain TM and short cytoplasmic domain 4B, SEMA4B, SEMAC, Semaphorin 4B, SemC, KIAA1745, MGC131831, SEM4B\_HUMAN.

Background: Semaphorins are a family of cell surface and secreted proteins that are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular semaphorin? domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. The transmembrane semaphorins are characterized by an additional 80 amino acid transmembrane domain and an 80-110 amino acid cytoplasmic domain. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA4B (semaphorin-4B), also known as SemC or SEMAC, is an 832 amino acid single-pass type I membrane protein that belongs to the semaphorin family and exists as two alternatively spliced isoforms. Containing one Ig-like C2-type (immunoglobulin-like) domain, a PSI domain and a single sema domain, SEMA4B is encoded by a gene located on human chromosome 15.

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months