



[Go to Product page](#)

Datasheet for ABIN1391541

## anti-SLK antibody (AA 885-931) (AbBy Fluor® 350)

### Overview

Quantity:	100 µL
Target:	SLK
Binding Specificity:	AA 885-931
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLK antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLK
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Sheep, Horse, Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	SLK
Alternative Name:	STK2 ( <a href="#">SLK Products</a> )
Background:	Synonyms: CTCL tumor antigen se20 9, LOSK, se20 9, Serine/threonine protein kinase 2, STK2,

## Target Details

---

SLK, SNF1 sucrose nonfermenting like kinase, STE20 like serine/threonine-protein kinase, STE20 related kinase, STK2, SLK\_HUMAN.

Background: SLK is a member of the serine/threonine kinase subfamily, Ste20. This subfamily is comprised of several mammalian kinases which exhibit sequence similarity to the *Saccharomyces cerevisiae* serine/threonine kinase Ste20, a protein involved in relaying signals from G protein-coupled receptors to cytosolic MAP kinase cascades. Members of this subfamily include KHS, GLK, YSK1, HPK1, Krs-1, Krs-2, GC kinase, HGK and SLK. SLK is a ubiquitously expressed protein that localizes to the cytoplasm and contains an N-terminal protein kinase domain, a central coiled-coil domain and a C-terminal ATH domain. SLK is activated through cleavage by caspase-3. SLK indirectly associates with microtubules and plays an important role in cellular stress, cell motility, cell death and cytoskeletal dynamics.

UniProt: [Q9H2G2](#)

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months