



[Go to Product page](#)

Datasheet for ABIN1391587
anti-UNC5B antibody (AA 188-260) (Biotin)

Overview

Quantity:	100 µL
Target:	UNC5B
Binding Specificity:	AA 188-260
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UNC5B antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UNC5B
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UNC5B
Alternative Name:	UNC5B/UNC5H2 (UNC5B Products)

Target Details

Background:	<p>Synonyms: Netrin receptor UNC5B, Netrin receptor UNC5B Precursor, p53 regulated receptor for death and le, p53 regulated receptor for death and le protein 1, p53-regulated receptor for death and le protein 1, p53RDL1, Protein unc-5 homolog 2, Protein unc-5 homolog B, Protein XUNC 5, Transmembrane receptor Unc5H2, Unc 5 homolog 2, Unc 5 homolog B, UNC 5B, Unc5 C.elegans homolog b, Unc5b, UNC5B_HUMAN, UNC5H2.</p> <p>Background: The UNC5H family of proteins act as transmembrane receptors for netrin-1 and play a crucial role in axon guidance and migration of neural cells. In fact, UNC5H receptors express widely in cells that migrate, where they bind the G protein G Alpha 1-2 to inhibit G protein signaling. Additionally, UNC5H receptors induce apoptosis when cleaved by a caspase, producing an intracellular fragment containing a death domain, but this activity is blocked by the binding of netrin-1. The expression of UNC5H receptors is down-regulated in multiple cancers, including colorectal, breast, ovary, uterus, stomach, lung, and kidney cancers. Hence, in the absence of netrin-1, UNC5H receptors act as tumor suppressors by inhibiting anchorage-independent growth and invasion, but mutation of these receptors provides a potential mechanism for tumorigenicity. UNC5H2, also designated unc-5 homolog B or p53-regulated receptor for death and life protein 1 (p53RDL1) is highly expressed in brain with lower levels of expression observed in developing lung, cartilage, kidney and hematopoietic and immune tissues.</p>
-------------	--

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months