antibodies -online.com





anti-UNC5D antibody (AA 112-200) (Biotin)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	UNC5D	
Binding Specificity:	AA 112-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UNC5D antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UNC5H4
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UNC5D
Alternative Name: UNC5D/UNC5H4 (UNC5D Products)	
Background: Synonyms: Netrin receptor UNC5D, Netrin receptor Unc5h4, Protein unc 5 homol	

homolog 4, Unc 5 homolog D C. elegans, Unc 5 homolog D, UNC 5D, Unc5h4, FLJ16019, KIAA1777.

Background: The UNC5H family of proteins act as transmembrane receptors for netrin-1 and play a crucial role in axon guidance and migration of neural cells. Additionally, when cleaved by a caspase to produce an intracellular fragment containing a death domain, UNC5H receptors induce apoptosis. This activity is blocked by the binding of netrin-1. In the absence of netrin-1, UNC5H receptors act as tumor suppressors by inhibiting anchorage-independent growth and invasion, but mutation of these receptors provides a potential mechanism for tumorigenicity. The expression of UNC5H receptors is down-regulated in multiple carcinomas, including colorectal, breast, ovary, uterus, stomach, lung, and kidney cancers. UNC5H4, also known as UNC5D (unc-5 homolog D), is single-pass type I membrane protein that is a member of the UNC5H netrin receptor family. Two isoforms of UNC5H4 exist due to alternative splicing events.

Application Details

Application Notes:

Expiry Date:

WB 1:300-5000

12 months

• •	
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.