antibodies -online.com





anti-BBS4 antibody (AA 431-519) (Alexa Fluor 350)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	BBS4
Binding Specificity:	AA 431-519
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BBS4 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BBS4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	BBS4
Alternative Name:	BBS4 (BBS4 Products)

Target Details

Background:

Synonyms: Bardet Biedl syndrome 4 protein, Bardet-Biedl syndrome 4 protein, Bbs4, BBS4_HUMAN.

Background: Bardet-Biedl syndrome (BBS) is a pleiotropic genetic disorder characterized by obesity, photoreceptor degeneration, polydactyly, hypogenitalism, renal abnormalities, and developmental delay. Other associated clinical findings in BBS patients include diabetes, hypertension, and congenital heart defects. BBS is a heterogeneous disorder, BBS genes map to eight genetic loci and encode eight proteins, BBS1-BBS8. Five BBS genes encode basal body or cilia proteins, suggesting that BBS is a ciliary dysfunction disorder. BBS4 is expressed in the olfactory epithelium and localizes to the centriolar satellites of centrosomes and basal bodies of primary cilia. BBS4 regulates the p150 subunit of the dynein transport machinery (DCTN1) to attract pericentriolar material-1 protein (PCM1) and its associated components to the satellites. Loss of BBS4 is correlated with obesity caused by abnormal lipid profiles, liver dysfunction, elevated insulin, and abnormal leptin levels.

Pathways:

Hedgehog Signaling, Tube Formation, Maintenance of Protein Location

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months