antibodies -online.com





anti-BBS9 antibody (AA 244-320) (Alexa Fluor 555)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | BBS9 |
| Binding Specificity: | AA 244-320 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BBS9 antibody is conjugated to Alexa Fluor 555 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human BBS9 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Dog,Sheep,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | BBS9 |
|-------------------|----------------------|
| Alternative Name: | BBS9 (BBS9 Products) |

Target Details

| - | | |
|---------------------|--|--|
| Background: | Synonyms: B1, D1, C18, PTHB1, Protein PTHB1, Bardet-Biedl syndrome 9 protein, Parathyroid | |
| | hormone-responsive B1 gene protein, BBS9 | |
| | Background: The BBSome complex is thought to function as a coat complex required for | |
| | sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for | |
| | ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is | |
| | mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and | |
| | contacts the BBSome. Rab8(GTP) enters the primary cilium and promotes extension of the | |
| | ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to | |
| | RAB3IP/Rabin8, the guanosyl exchange factor (GEF) for Rab8 and then the Rab8-GTP localizes | |
| | to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary | |
| | membrane. Required for proper BBSome complex assembly and its ciliary localization. | |
| Gene ID: | 27241 | |
| UniProt: | Q3SYG4 | |
| Pathways: | Hedgehog Signaling | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| | | |

| | | 1 . | |
|---|-------|-------|---|
| ш | land | linc | ٩ |
| | ıanıu | 11111 | J |
| | | | |

Expiry Date:

12 months