# antibodies .- online.com





Datasheet for ABIN1391861

# anti-CDKL5 antibody (AA 401-500) (Alexa Fluor 555)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	CDKL5	
Binding Specificity:	AA 401-500	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CDKL5 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CDKL5	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	CDKL5
Alternative Name:	CDKL5 (CDKL5 Products)
Background:	Synonyms: Cdkl5, CDKL5_HUMAN, Cyclin dependent kinase 5 transcript, Cyclin-dependent

kinase-like 5, EIEE2, ISSX, Serine/threonine kinase 9, Serine/threonine-protein kinase 9, Stk9. Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin-dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. CDKL5 (cyclin-dependent kinase-like 5) is a 1030 amino acid protein that belongs to the CMGC Ser/Thr protein kinase family. Expressed in brain, lung, kidney, prostate, ovary, placenta, pancreas and testis, CDKL5 is thought to play a role in cell cycle regulation. Defects in CDKL5 are a cause of several disorders, such as X-linked infantile spasm syndrome and Rett syndrome.

Pathways:

Regulation of Cell Size

### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	