



[Go to Product page](#)

Datasheet for ABIN1391947

anti-EN2 antibody (AA 281-333) (Biotin)

1 Publication

Overview

Quantity:	100 µL
Target:	EN2
Binding Specificity:	AA 281-333
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EN2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EN2/Engrailed 2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Sheep,Pig
Purification:	Purified by Protein A.

Target Details

Target:	EN2
Alternative Name:	EN2/Engrailed 2 (EN2 Products)

Target Details

Background:	<p>Synonyms: Homeobox protein engrailed-2, Homeobox protein en-2, Hu-En-2, EN2</p> <p>Background: The engrailed-2 gene, EN2, a murine homolog of the Drosophila homeobox gene engrailed (EN), is required for midbrain and cerebellum development and dorsal/ventral patterning of the limbs as well as apical ectodermal ridge formation. In Drosophila, the EN gene plays an important role during development in segmentation, where it is required for the formation of posterior compartments. Human EN-1 and EN-2 are homeodomain-containing proteins and have been implicated in the control of pattern formation during development of the central nervous system. Different mutations in the mouse homologs, EN-1 and EN-2, produce different developmental defects that frequently are lethal. EN-1 is highly expressed by essentially all dopaminergic neurons in the substantia nigra and ventral tegmentum. EN-1 and EN-2 regulate expression of a-synuclein, a gene that is genetically linked to Parkinson's disease. During early brain development mouse EN-2 is expressed in a broad band across most of the mid-hindbrain region. EN-2 is also expressed in mouse myoblasts and has been associated with cerebellar hypoplasia.</p>
-------------	--

Gene ID:	2020
----------	------

UniProt:	P19622
----------	------------------------

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	--

Preservative:	ProClin
---------------	---------

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

Storage:	-20 °C
----------	--------

Handling

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months

Publications

Product cited in: De Paoli, Gogalic, Sauer, Preininger, Pandha, Simpson, Horvath, Marquette: "Multiplatform Biomarker Discovery for Bladder Cancer Recurrence Diagnosis." in: **Disease markers**, Vol. 2016, pp. 4591910, (2016) ([PubMed](#)).