-online.com antibodies

# Datasheet for ABIN1392077 anti-CDK17 antibody (AA 301-400) (Alexa Fluor 555)



Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | CDK17  |
| Binding Specificity: | AA 301-400   |
| Reactivity:          | Sheep  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This CDK17 antibody is conjugated to Alexa Fluor 555   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

# Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human PCTAIRE2 |
|-----------------------|--|
| lsotype:              | lgG  |
| Cross-Reactivity:     | Sheep  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Chicken,Rabbit                 |
| Purification:         | Purified by Protein A.                                       |
| Target Details        |  |
| Target:               | CDK17  |
| Alternative Name:     | PCTAIRE2 (CDK17 Products)                                    |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1392077 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

## Target Details

#### Background:

Synonyms: PCTAIRE-2, PCTAIRE-mot protein kinase 2, PCTK2, Protein kinase cdc2-related PCTAIRE-2, Serine/threonine-protein kinase PCTAIRE-2, 6430598J10Rik, EC 2.7.1.37, MGC25109, CDK17\_HUMAN.

Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with cyclins to phosphorylate key substrates involved in cell cycle progression. Another family of proteins, Cdk inhibitors, also play a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Members of the Cdk family include Cdk2Cdk8, PCTAIRE-13, PITALRE and PITSLRE. PCTAIRE-1, PCTAIRE-2 and PCTAIRE-3 comprise a subfamily of cdc2-related serine/threonine kinases. PCTAIRE-1, which is expressed primarily in mammalian brain, interacts with a variety of proteins and is thought to be part of a multiple signal transduction cascade. PCTAIRE-2, also expressed in brain, may be important in terminally differentiated neurons.

# Application Details

| Application Notes: | IF(IHC-P) 1:50-200    |
|--------------------|-----------------------|
|                    | IF(IHC-F) 1:50-200    |
|                    | IF(ICC) 1:50-200      |
| Restrictions:      | For Research Use only |

### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1392077 | 03/08/2024 | Copyright antibodies-online. All rights reserved.