Datasheet for ABIN1392079 anti-CDK17 antibody (AA 301-400) (Biotin)

-online.com antibodies



\sim							
()	\/		rv	Т.		۱۸/	I
U	v	C	I V	L.	C	vv	

Quantity:	100 µL
Target:	CDK17
Binding Specificity:	AA 301-400
Reactivity:	Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK17 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PCTAIRE2
lsotype:	IgG
Cross-Reactivity:	Sheep
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Chicken,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	CDK17
Alternative Name:	PCTAIRE2 (CDK17 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1392079 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:

Synonyms: PCTAIRE-2, PCTAIRE-mot protein kinase 2, PCTK2, Protein kinase cdc2-related PCTAIRE-2, Serine/threonine-protein kinase PCTAIRE-2, 6430598J10Rik, EC 2.7.1.37, MGC25109, CDK17_HUMAN.

Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with cyclins to phosphorylate key substrates involved in cell cycle progression. Another family of proteins, Cdk inhibitors, also play a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Members of the Cdk family include Cdk2Cdk8, PCTAIRE-13, PITALRE and PITSLRE. PCTAIRE-1, PCTAIRE-2 and PCTAIRE-3 comprise a subfamily of cdc2-related serine/threonine kinases. PCTAIRE-1, which is expressed primarily in mammalian brain, interacts with a variety of proteins and is thought to be part of a multiple signal transduction cascade. PCTAIRE-2, also expressed in brain, may be important in terminally differentiated neurons.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months