# antibodies .- online.com





# anti-GPRC5A antibody (AA 1-33) (Alexa Fluor 488)



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL	
Target:	GPRC5A	
Binding Specificity:	AA 1-33	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GPRC5A antibody is conjugated to Alexa Fluor 488	
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RAI3	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	GPRC5A	
Alternative Name:	RAI3 (GPRC5A Products)	
Background:	ackground: Synonyms: G protein coupled receptor family C group 5 member A, G-protein coupled rec	

family C group 5 member A, GPCR 5A, GPCR5A, Gprc 5a, Gprc5a, Orphan G protein coupling receptor PEIG 1, Orphan G protein coupling receptor PEIG1, Orphan G-protein-coupling receptor PEIG-1, RAI 3, RAI3\_HUMAN, Raig 1, RAIG-1, Raig1, Retinoic acid induced 3, Retinoic acid induced gene 1 protein, Retinoic acid induced protein 3, Retinoic acid responsive, Retinoic acid responsive gene, Retinoic acid-induced gene 1 protein, Retinoic acid-induced protein 3.

Background: RAI3 is an transmembrane G-protein-coupled receptor that affects many essential biological processes including embryogenesis, cell growth, differentiation, and apoptosis. RAI3 may also be involved in maintaining homeostasis of epithelial cells. Retinoic acid receptors directly regulate RAI3 during its transcription in embryonal carcinoma differentiation. RAI3 expression is upregulated in most tumor cell lines that express mutant p53, suggesting that p53 interacts with the promoter of RAI3 and represses its expression at the beginning of apoptosis. RAI3 is a potential molecular target for diagnosing breast cancer, and selective suppression of signals from RAI3 may have a place in breast cancer treatments.

## **Application Details**

Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months