



[Go to Product page](#)

Datasheet for ABIN1392204

anti-NOMO1 antibody (AA 701-800) (Alexa Fluor 647)

Overview

Quantity:	100 µL
Target:	NOMO1
Binding Specificity:	AA 701-800
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOMO1 antibody is conjugated to Alexa Fluor 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NOMO1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	NOMO1
Alternative Name:	NOMO1 (NOMO1 Products)
Background:	Synonyms: Nodal modulator 1, NOMO 1, Nomo, NOMO1, NOMO1_HUMAN, PM5, pM5 protein,

Target Details

pM5 protein telomeric copy, NOMO1_HUMAN.

Background: Three highly similar proteins termed NOMO1, NOMO2 and NOMO3, are encoded by a gene mapping to a region of duplication on the p arm of human chromosome 16. All three NOMO proteins share similar functions and have been difficult to characterize individually. NOMO1 (Nodal modulator 1), also known as PM5, is a 1,222 amino acid highly conserved single-pass type I membrane protein expressed in colon tumor tissue and normal colonic mucosa. NOMO proteins are novel antagonists of Nodal signaling which interact with Nicalin to form a Nicalin-NOMO complex, and are rapidly degraded or stabilized by Nicalin. NOMO proteins were once considered candidates for the development of pseudoxanthoma elasticum (PXE), a heritable disorder of connective tissue, as the NOMO genes are located in close proximity to the gene responsible for PXE development (MRP6).

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months