



[Go to Product page](#)

Datasheet for ABIN1392211
anti-OTX1 antibody (AA 1-100) (Biotin)

Overview

Quantity:	100 µL
Target:	OTX1
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OTX1 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human OTX1
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	OTX1
Alternative Name:	OTX1 (OTX1 Products)
Background:	Synonyms: Homeobox protein OTX 1, Homeobox protein OTX1, MGC15736, Orthodenticle 1,

Target Details

Orthodenticle homeobox 1, Orthodenticle homolog 1, Orthodenticle1, Otx 1, otx1, OTX1_HUMAN.

Background: Transcription factors, OTX1 and OTX2, are two murine homologs of the *Drosophila* orthodenticle (OTD), show a limited amino acid sequence divergence. OTX1 and OTX2 play an important role during early and later events required for proper brain development in that they are involved in the processes of induction, specification and regionalization of the brain. OTX1 is involved in corticogenesis, sensory organ development and pituitary functions, while OTX2 is necessary earlier in development, for the correct anterior neural plate specification and organization of the primitive streak. OTX2 is also required in the early specification of the neuroectoderm, which is destined to become the fore-midbrain, and both OTX1 and OTX2 co-operate in patterning the developing brain through a dosage-dependent mechanism. A molecular mechanism depending on a precise threshold of OTX proteins is necessary for the correct positioning of the isthmus region and for anterior brain patterning. The genes which encode OTX1 and OTX2 map to human chromosomes 2p15 and 14q21-q22, respectively.

Application Details

Application Notes: IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months