# antibodies -online.com





## anti-ZIC2 antibody (AA 201-300) (Biotin)



_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μL	
Target:	ZIC2	
Binding Specificity:	AA 201-300	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZIC2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Zic2	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Chicken	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	ZIC2
Alternative Name:	Zic2 (ZIC2 Products)

#### **Target Details**

Storage Comment:

Expiry Date:

l arget Details	
Background:	Synonyms: HPE 5, HPE5, Odd paired homolog Drosophila, Zic 2, Zic family member 2 odd
	paired Drosophila homolog, Zic family member 2, ZIC2, ZIC2_HUMAN, Zinc finger protein of the
	cerebellum 2, Zinc finger protein ZIC 2, Zinc finger protein Zic2.
	Background: Zic2 is a C2H2 zinc finger transcription factor that influences forebrain
	development. Zic2 is a transcriptional repressor and may regulate tissue specific expression of
	dopamine receptor D1. Zic2 transcript is abundant in the dorsal neural tube/spinal cord, and in
	the hindbrain. A polyhistidine tract polymorphism in this gene may be associated with
	increased risk of neural tube defects. This gene is closely linked to a gene encoding zinc finger
	protein of the cerebellum 5, a related family member on chromosome 13.
Pathways:	Tube Formation
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C for 12 months.

12 months