# antibodies -online.com





## anti-TAS2R16 antibody (FITC)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	TAS2R16
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAS2R16 antibody is conjugated to FITC
Application:	Western Blotting (WB)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TAS2R16	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:

TAS2R16

3 - 1	
Alternative Name:	Tas2r16 (TAS2R16 Products)
Background:	Synonyms: TAS2R-16, T2R16, Taste receptor type 2 member 16, T2R16_HUMAN.
	Background: T2R16 is a 291 amino acid multi-pass membrane protein that belongs to the G-
	protein coupled receptor T2R family. Acting in the oral cavity and the gastrointestinal tract,
	T2R16 is a gustducin-coupled receptor that is implicated in the perception of bitter compounds.

T2R16 mediates responses to certain taste through PLC 2, a phospholipase C selectively expressed in taste tissue, and the calcium-regulated cation channel TRPM5. Expressed in a subset of gustducin-positive taste receptor cells of the tongue and epithelia, T2R16 confers bitter perception of salicin to non-taster mice. The gene that encodes T2R16 consists of 996 bases and maps to human chromosome 7q31. Chromosome 7 houses over 1,000 genes, comprises nearly 5 % of the human genome and has been linked to Osteogenesis imperfecta, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome. The Lys-172 polymorphism in T2R16 is associated with genetic susceptibility to alcoholism.

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months