# antibodies -online.com





# anti-PKD1L3 antibody (AA 121-220) (FITC)



Go to Product page

$\sim$				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	PKD1L3
Binding Specificity:	AA 121-220
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKD1L3 antibody is conjugated to FITC
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PKD1L3
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

## **Target Details**

Target:	PKD1L3
Alternative Name:	PKD1L3 (PKD1L3 Products)
Background: Synonyms: PC1 like 3 protein, Polycystic kidney disease 1 like 3, Polycystic kidney disease	

protein 1 like 3, Polycystin 1 like 3, Polycystin 1L3, PK1L3\_HUMAN.

Background: Polycystin-1L3 is a 1,732 amino acid multi-pass membrane protein that contains one PLAT domain, one GPS domain and one C-type lectin domain. Expressed at high levels in placenta and present at lower levels in lung and heart, Polycystin-1L3 is thought to function as an ion-channel regulator that may interact with Polycystin-L and play a role in heteromeric taste channels. The gene encoding Polycystin-1L3 maps to human chromosome 16, which encodes over 900 genes and comprises nearly 3 % of the human genome. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is also associated with chromosome 16, as is Crohn's disease, which is a gastrointestinal inflammatory condition.

## **Application Details**

IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months