antibodies -online.com







anti-PIRT antibody (AA 51-137) (Alexa Fluor 350)



| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | PIRT | |
| Binding Specificity: | AA 51-137 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PIRT antibody is conjugated to Alexa Fluor 350 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PIRT | |
|---|--|--|
| Isotype: | IgG | |
| Predicted Reactivity: Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Rabbit | | |
| Purification: | Purified by Protein A. | |

Target Details

| - anger pleame | | |
|-------------------|---|--|
| Target: | PIRT | |
| Alternative Name: | PIRT (PIRT Products) | |
| Background: | Synonyms: hCG 1776018. Phosphoinositide interacting regulator of transient receptor potential | |

channels, Phosphoinositide-interacting protein, Pirt, PIRT_HUMAN.

Background: PIRT is a 137 amino acid multi-pass membrane protein. Highly conserved among vertebrates, PIRT consists of two transmembrane domains and one putative C-terminal phosphoinositide-binding domain. Although PIRT is expressed in peripheral nervous system, with highest levels in dorsal root ganglion and trigeminal neurons, and lowest levels in sympathetic and enteric neurons, it is not expressed in spinal cord. PIRT is a required component of the VR1 complex, which positively regulates VR1, a sensor of both noxious heat and capsaicin. Correspondingly, PIRT knockout results in impaired responses to noxious heat and capsaicin exposure, while VR1 remains unaltered. The gene that encodes PIRT maps to human chromosome 17p13.1.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |