



[Go to Product page](#)

Datasheet for ABIN1392452
anti-APBB3 antibody (AA 401-486) (FITC)

Overview

Quantity:	100 µL
Target:	APBB3
Binding Specificity:	AA 401-486
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APBB3 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human APBB3/FE65L2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep
Purification:	Purified by Protein A.

Target Details

Target:	APBB3
Alternative Name:	APBB3/FE65L2 (APBB3 Products)

Target Details

Background: Synonyms: Amyloid beta A4 precursor protein binding family B member 3, Amyloid beta A4 precursor protein-binding family B member 3, amyloid beta precursor protein binding family B member 3, amyloid precursor interacting protein, Apbb3, APBB3_HUMAN, Fe65 like protein 2, FE65L2, Protein Fe65-like 2, SRA.

Background: Fe65L2 is a 486 amino acid protein that contains one WW domain and two PID domains. Binding to the intracellular domain of the -Amyloid precursor protein, Fe65L2 is thought to modulate the internalization and, therefore, the accessibility and function of -Amyloid. Via its ability to control the intracellular accumulation of -Amyloid, Fe65L2 is thought to play a role in the pathogenesis of Alzheimer's disease. Fe65L2 exists as four alternatively spliced isoforms designated isoform I, isoform II, isoform III and isoform IV. Fe65L2 interacts with Amyloid-like protein and is encoded by a gene located on human chromosome 5, which contains 181 million base pairs and comprises nearly 6 % of the human genome. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

Gene ID: 10307

UniProt: [095704](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months