# antibodies -online.com





## anti-LZIC antibody (AA 31-130) (FITC)



	1
Overview	/

Quantity:100 μLTarget:LZICBinding Specificity:AA 31-130Reactivity:MouseHost:RabbitClonality:PolyclonalConjugate:This LZIC antibody is conjugated to FITCApplication:Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence		
Binding Specificity: AA 31-130  Reactivity: Mouse  Host: Rabbit  Clonality: Polyclonal  Conjugate: This LZIC antibody is conjugated to FITC	Quantity:	100 μL
Reactivity: Mouse  Host: Rabbit  Clonality: Polyclonal  Conjugate: This LZIC antibody is conjugated to FITC	Target:	LZIC
Host: Rabbit Clonality: Polyclonal Conjugate: This LZIC antibody is conjugated to FITC	Binding Specificity:	AA 31-130
Clonality: Polyclonal  Conjugate: This LZIC antibody is conjugated to FITC	Reactivity:	Mouse
Conjugate: This LZIC antibody is conjugated to FITC	Host:	Rabbit
	Clonality:	Polyclonal
Application: Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluores	Conjugate:	This LZIC antibody is conjugated to FITC
(Paraffin-embedded Sections) (IF (p))	Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human LZIC
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	LZIC
Alternative Name:	LZIC (LZIC Products)

#### **Target Details**

#### Background:

Synonyms: Leucine zipper and CTNNBIP1 domain containing protein, Leucine zipper and ICAT homologous domain containing protein, leucine zipper domain and ICAT homologous domain containing, LZIC antibodyprotein LZIC, LZIC\_HUMAN.

Background: LZIC is a 190 amino acid protein that belongs to the CTNNBIP1 family. Ubiquitously expressed, LZIC is found at highest levels in kidney and is upregulated in a few gastric carcinomas. The gene encoding LZIC maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8 % of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson?s disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

#### **Application Details**

/\nn	lication	MOTAC:
$\neg$ uu	IICatiOII	INULES

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months