# antibodies -online.com





# anti-Calsyntenin 1 antibody (AA 501-600) (Alexa Fluor 350)



Go to Product page

( )	۱ ۱	$\cap$	r	/1	$\cap$	۱ ۸	1
0	'V	ㄷ	I١	νı	$\Box$	V	۷

Quantity:	100 μL	
Target:	Calsyntenin 1 (CLSTN1)	
Binding Specificity:	AA 501-600	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Calsyntenin 1 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Calsyntenin 1	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Pig,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	Calsyntenin 1 (CLSTN1)
Alternative Name:	CLSTN1/Calsyntenin 1 (CLSTN1 Products)
Background:	Synonyms: Alc alpha, Alc-alpha, Alcadein alpha 1, Alcadein alpha, Alcadein-alpha, alcalpha1,

alcalpha2, Alzheimer related cadherin like protein, Alzheimer-related cadherin-like protein, Calsyntenin-1, Calsyntenin-1, CLSTN 1, Clstn1, CS1, CSTN1, CSTN1\_HUMAN, CTF1-alpha, FLJ32258, KIAA0911, Non classical cadherin XB31alpha, Non-classical cadherin XB31alpha, PIK3CD, SAlc-alpha, XB31alpha.

Background: Members of the calsyntenin protein family are localized to the post-synaptic membrane of exicitatory central nervous system (CNS) synapses. Calsyntenin-1, also known as CSTN1, PIK3CD, Alzheimer-related cadherin-like protein, non-classical cadherin XB31alpha, KIAA0911, ALC-ALPHA, alcalpha1, alcalpha2 or FLJ32258, is a 981 amino acid single-pass type I membrane protein that localizes to the membrane of endoplasmic reticulum, Golgi apparatus, cell projections and postsynaptic cells. Expressed in brain, calsyntenin-1 is also found at lower levels in placenta, skeletal muscle, heart and kidney. Calsyntenin-1 binds synaptic Ca2+ with its cytoplasmic domain and plays a role in extracellular proteolysis. Calsyntenin-1 is also known to form a complex with X11 Beta and APP to suppress the metabolic cleavage of APP, and docks vesicular cargo to KLC1. Calsyntenin-1 may be related to the development or progression of Alzheimer?s disease, and two calsyntenin-1 isoforms are produced as a result of alternative splicing events.

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months