# antibodies -online.com





## anti-SH3PXD2A antibody (AA 261-360) (FITC)



| $\sim$  |          |
|---------|----------|
| ( )\/\  | rview    |
| $\circ$ | 1 410 44 |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | SH3PXD2A  |
| Binding Specificity: | AA 261-360  |
| Reactivity:          | Rat   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SH3PXD2A antibody is conjugated to FITC  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FISH |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Rat  |
| Predicted Reactivity: | Human, Mouse   |
| Purification:         | Purified by Protein A.                                   |

### Target Details

| Target:           | SH3PXD2A                 |
|-------------------|--------------------------|
| Alternative Name: | FISH (SH3PXD2A Products) |

#### **Target Details**

#### Background:

Synonyms: Adapter protein TKS5, Five SH3 domain-containing protein, SH3 and PX domain-containing protein 2A, SH3 multiple domains protein 1, Sh3md1, Sh3pxd2a, SPD2A\_HUMAN, TKs5, Tyrosine kinase substrate with five SH3 domains.

Background: Fish, a potential Src substrate, is a broadly expressed adaptor protein containing five SH3 domains and a phox homology (PX) domain (1). The Src family of protein tyrosine kinases act in signal transduction pathways (2-4). Src kinases vary in expression but are strongly regulated in vivo, catalytic activity is repressed by interacting with the SH3 domain (5-7). In Src-transformed fibroblasts and in normal cells treated with certain growth factors fish is tyrosine-phosphorylated (1). Treatment of cells with cytochalasin D results in rapid tyrosine phosphorylation of fish, along with activation of Src (1). Fish is likely to be involved in tyrosine kinase signaling and may have a role in cytoskeletal changes (1).

#### **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |