antibodies -online.com





anti-PICALM antibody (AA 251-350) (Alexa Fluor 350)



Go to Product page

\sim			
	N/P	r\/I	$\Theta \backslash \Lambda /$

Quantity:	100 μL
Target:	PICALM
Binding Specificity:	AA 251-350
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PICALM antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PICALM/CALM
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Horse,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	PICALM
Alternative Name:	PICALM/CALM (PICALM Products)

Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		\sim	<i>.</i>	ч.

Synonyms: Clathrin assembly lymphoid myeloid leukemia, Clathrin assembly lymphoid myeloid leukemia protein, CLTH, LAP, Phosphatidylinositol-binding clathrin assembly protein, PICAL_HUMAN, PICALM.

Background: Clathrin-coated pits and vesicles are assembled for receptor-mediated endocytosis through interaction with clathrin associated protein complexes. Vesicle transport is mediated from the trans-Golgi network by the adapter complex AP-1 and from the plasma membrane by the AP-2 complex. The AP-1 and AP-2 adapter protein complexes consist of clathrin binding adaptin proteins and two smaller subunits known as AP50 and AP17. The Alpha- and Beta-Adaptin chains have a similar two-domain organization with C-terminal domains that vary in both sequence and length. Alpha-Adaptin splice variants A and C display variable relative expression levels and differential distribution in different tissues. AP180 (also designated AP-3 or F1-20) is a synapse-specific clathrin assembly protein. The protein CALM (clathrin assembly protein lymphoid myeloid leukemia) is highly homologous to AP180 and may also be involved in clathrin assembly.

Pathways:

Synaptic Membrane

Application Details

Δ nn	lication	Notae.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months