antibodies -online.com





anti-PICALM antibody (AA 251-350) (Biotin)



| () | ve | K\ / | | A . |
|-----|--------|------|---|-----|
| | \cup | 1 V/ | - | V۷ |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | PICALM |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PICALM antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PICALM/CALM |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Horse,Zebrafish |
| Purification: | Purified by Protein A. |

Target Details

| Target: | PICALM |
|-------------------|-------------------------------|
| Alternative Name: | PICALM/CALM (PICALM Products) |

Target Details

Background:

Synonyms: Clathrin assembly lymphoid myeloid leukemia, Clathrin assembly lymphoid myeloid leukemia protein, CLTH, LAP, Phosphatidylinositol-binding clathrin assembly protein, PICAL_HUMAN, PICALM.

Background: Clathrin-coated pits and vesicles are assembled for receptor-mediated endocytosis through interaction with clathrin associated protein complexes. Vesicle transport is mediated from the trans-Golgi network by the adapter complex AP-1 and from the plasma membrane by the AP-2 complex. The AP-1 and AP-2 adapter protein complexes consist of clathrin binding adaptin proteins and two smaller subunits known as AP50 and AP17. The Alpha- and Beta-Adaptin chains have a similar two-domain organization with C-terminal domains that vary in both sequence and length. Alpha-Adaptin splice variants A and C display variable relative expression levels and differential distribution in different tissues. AP180 (also designated AP-3 or F1-20) is a synapse-specific clathrin assembly protein. The protein CALM (clathrin assembly protein lymphoid myeloid leukemia) is highly homologous to AP180 and may also be involved in clathrin assembly.

Pathways:

Synaptic Membrane

Application Details

| Λnn | lication | Motac. |
|--------|----------|---------|
| \neg | псанон | TAUTES. |

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |

| | | 1. | |
|----------|-------|-------|---|
| \vdash | land | ling | |
| 1 1 | ıaııu | 11111 | ۰ |
| | | | |

Expiry Date:

12 months