# antibodies -online.com





## anti-Tppp antibody (AA 61-150) (Biotin)



Go to Product page

( )	1/0	r\ / I	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Тррр	
Binding Specificity:	AA 61-150	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Tppp antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TPPP
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Chicken
Purification:	Purified by Protein A.

### **Target Details**

Target:	Тррр
Alternative Name:	TPPP (Tppp Products)

#### **Target Details**

#### Background:

Synonyms: Tubulin Polymerization Promoting Protein 24, 25 kDa brain specic protein, 25 kDa brain-specic protein, Brain specic protein p25 alpha, Glycogen synthase kinase 3 GSK3 inhibitor p24, p24, p25, p25-alpha, p25alpha, TPPP, TPPP/p25, TPPP\_HUMAN, TPPP1, Tubulin polymerization promoting protein, Tubulin polymerization-promoting protein.

Background: Tubulin family members are globular proteins important in the assembly of microtubules. Microtubules are structural components that play important roles in mitosis, cytokinesis and vesicle transport. TPPP (Tubulin polymerization-promoting protein), also known as p24 and p25, is a widely expressed 219 amino acid protein found in the perinuclear region of the cytoplasm. TPPP may form dimers and functions in polymerizing tubulin into double-walled tubules, polymorphic aggregates, or stabilized blocks. TPPP overexpression prevents formation of the mitotic spindle assembly and breakdown of the nuclear envelope. TPPP is phosphorylated by TPK II and is encoded by a gene that maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6 % of the human genome.

#### **Application Details**

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months