



[Go to Product page](#)

Datasheet for ABIN1392789

anti-HAAO antibody (AA 101-200) (AbBy Fluor® 350)

Overview

Quantity:	100 µL
Target:	HAAO
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HAAO antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HAAO
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HAAO
Alternative Name:	HAAO (HAAO Products)

Target Details

Background:	<p>Synonyms: 3 HAO,3 hydroxyanthranilate 3 4 dioxygenase,3 hydroxyanthranilate dioxygenase,3 hydroxyanthranilate oxygenase,3 hydroxyanthranilic acid dioxygenase,3-HAO,3-hydroxyanthranilate 3,3-hydroxyanthranilate 3,4-dioxygenase,3-hydroxyanthranilate oxygenase,3-hydroxyanthranilic acid dioxygenase,3HAO_HUMAN,4-dioxygenase,haao,HAD,HAO.</p> <p>Background: The 3-HAO family of proteins consists of intramolecular dioxygenases that contain non-heme ferrous iron. HAAO (3-hydroxyanthranilate 3,4-dioxygenase), also known as HAO or 3-HAO, is a 286 amino acid monomeric cytoplasmic protein that belongs to the 3-HAO family. Utilizing Fe²⁺ ions as a cofactor, HAAO participates in the formation of quinolinic acid (QUIN), and excitotoxin, by catalyzing the oxidative ring opening of 3-hydroxyanthranilate to 2-amino-3-carboxymuconate semialdehyde. HAAO may play a role in disorders that are associated with varying levels of QUIN expression, and is widely expressed in peripheral organs, including kidney and liver. Existing as two alternatively spliced isoforms, the gene encoding HAAO maps to human chromosome 2p21 and mouse chromosome 17 E4.</p>
-------------	--

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months