



[Go to Product page](#)

Datasheet for ABIN1392901
anti-TBCB antibody (AA 1-100) (Biotin)

Overview

Quantity:	100 µL
Target:	TBCB
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBCB antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CKAP1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	TBCB
Alternative Name:	CKAP1/TBCB (TBCB Products)
Background:	Synonyms: CG22, CKAPI, Cytoskeleton associated protein 1, Cytoskeleton associated protein

Target Details

CKAPI, Cytoskeleton-associated protein 1, Cytoskeleton-associated protein CKAPI, TBCB, TBCB_HUMAN, Tubulin folding cofactor B, Tubulin specific chaperone B, Tubulin-folding cofactor B, Tubulin-specific chaperone B.

Background: Microtubules, the primary component of the cytoskeletal network, are highly dynamic structures composed of Alpha/Beta Tubulin heterodimers. Biosynthesis of functional microtubules involve the participation of several chaperones, termed Tubulin folding cofactors A (TBCA), B (TBCB), D (TBCD), E (TBCE) and C (TBCC), that act on folding intermediates downstream of the cytosolic chaperon, alternatively named TCP. TBCB (tubulin folding cofactor B), also known as CG22, CKAP1 or CKAPI, is a 244 amino acid cytoplasmic protein containing one CAP-Gly domain and is widely expressed. TBCB is involved in the regulation of tubulin heterodimer dissociation and may function as a negative regulator of axonal growth.

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months