



[Go to Product page](#)

Datasheet for ABIN1392903

## anti-CLN5 antibody (AA 61-120) (Alexa Fluor 350)

### Overview

Quantity:	100 µL
Target:	CLN5
Binding Specificity:	AA 61-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLN5 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLN5
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	CLN5
Alternative Name:	CLN5 ( <a href="#">CLN5 Products</a> )

## Target Details

---

Background:	<p>Synonyms: Ceroid lipofuscinosis neuronal 5, Ceroid-lipofuscinosis neuronal protein 5, CLN5, CLN5_HUMAN, NCL, Protein CLN5.</p> <p>Background: Neuronal ceroid-lipofuscinose (NCL), also designated Batten disease, comprises a group of recessively inherited, progressive neurodegenerative diseases found in children. NCL is characterized by atrophy of the brain and an accumulation of lysosome derived fluorescent bodies found in many cells, especially neurons. Symptoms of NCL include a failure of psychomotor development, seizures, impaired vision and premature death. The eight genes/proteins associated with NCL are designated CLN1-CLN8. Mutations in six of these genes results in a distinct type of NCL-disease, the six genes/proteins are CLN1 (encoding PPT1, a protein thiolesterase), CLN2 (encoding the serine protease TPP1), CLN3, CLN5, CLN6 and CLN8. A single base duplication mutation in dog and cow CLN5 has been shown to cause NCL.</p>
-------------	--

## Application Details

---

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months