# antibodies -online.com





# anti-CLN5 antibody (AA 61-120) (Alexa Fluor 555)



( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Quantity:	100 μL
Target:	CLN5
Binding Specificity:	AA 61-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLN5 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLN5	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	CLN5
Alternative Name:	CLN5 (CLN5 Products)

#### **Target Details**

#### Background:

Synonyms: Ceroid lipofuscinosis neuronal 5, Ceroid-lipofuscinosis neuronal protein 5, CLN5, CLN5\_HUMAN, NCL, Protein CLN5.

Background: Neuronal ceroid-lipofuscinose (NCL), also designated Batten disease, comprises a group of recessively inherited, progressive neurodegenerative diseases found in children. NCL is characterized by atrophy of the brain and an accumulation of lysosome derived fluorescent bodies found in many cells, especially neurons. Symptoms of NCL include a failure of psychomotor development, seizures, impaired vision and premature death. The eight genes/proteins associated with NCL are designated CLN1-CLN8. Mutations in six of these genes results in a distinct type of NCL-disease, the six genes/proteins are CLN1 (encoding PPT1, a protein thiolesterase), CLN2 (encodeing the serine protease TPP1), CLN3, CLN5, CLN6 and CLN8. A single base duplication mutation in dog and cow CLN5 has been shown to cause NCL.

### **Application Details**

/\nn	lication	MOTAC:
$\neg$ uu	lication	INUICS.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months