antibodies -online.com





anti-NMNAT1 antibody (AbBy Fluor® 350)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	NMNAT1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NMNAT1 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Nmnat1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NMNAT1	
Alternative Name:	Nmnat1 (NMNAT1 Products)	
Background:	Synonyms: NaMN adenylyltransferase 1, Nicotinamide mononucleotide adenylyltransferase 1,	
	nicotinamide nucleotide adenylyltransferase 1, nicotinate nucleotide adenylyltransferase 1,	

Nicotinate-nucleotide adenylyltransferase 1, NMN adenylyltransferase 1, NMNA1_HUMAN, Nmnat 1, Nmnat1, Nmnat-1, OTTHUMP00000001731, OTTHUMP00000001732, OTTHUMP00000035892, PNAT 1, PNAT1, pyridine nucleotide adenylyltransferase 1.

Background: Nicotinamide adenine dinucleotide (NMNAT) is an essential cofactor involved in fundamental processes in cell metabolism. NMNAT plays a key role in NAD(+) biosynthesis, catalysing the condensation of nicotinamide mononucleotide and ATP, and yielding NAD(+) and pyrophosphate. NMNAT appears to be a substrate of nuclear kinases and contains at least three potential phosphorylation sites. The interaction of NMNAT with nuclear proteins is likely to be modulated by phosphorylation. NMNAT is widely expressed with highest levels in skeletal muscle, heart, liver and kidney.

Application Details

Application Notes:

Storage:

Expiry Date:

Storage Comment:

Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

IF(IHC-P) 1:50-200

-20 °C

12 months