

Datasheet for ABIN1393055

anti-DMPK antibody (AA 51-120) (AbBy Fluor® 555)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DMPK
Binding Specificity:	AA 51-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DMPK antibody is conjugated to AbBy Fluor® 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DMPK
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.

Target Details

Target:	DMPK
Alternative Name:	DMPK (DMPK Products)

Target Details

Background: Synonyms: Dystrophia myotonica protein kinase, DM 1, DM, DM kinase, DM protein kinase, DM-kinase, DM1, DM1 protein kinase, DM1PK, DMK, DMPK, DMPK_HUMAN, Dystrophia myotonica 1, Dystrophia myotonica protein kinase, MDPK, MT PK, MT-PK, Myotonic dystrophy associated protein kinase, Myotonic dystrophy protein kinase, Myotonin protein kinase A, Myotonin protein kinase, Myotonin-protein kinase, Thymopoietin homolog.

Background: Myotonic dystrophy protein kinase is a multi-domain protein kinase found in muscle that is activated in response to G protein second messengers and proteolysis (1). DMPK is implicated in myotonic muscular dystrophy (DM), an autosomal dominant-inherited disorder that predominately affects skeletal and cardiac muscle and causes defects in cardiac conduction (2,3). DM arises through expansion of CTG repeats in the 3'-UTR of the DMPK gene (4). Mutant DMPK transcripts with an extended region of CUG repeats are retained in the nucleus (5). These transcripts also influence the expression of the DM locus-associated homeodomain protein (DMAHP)/SIX5, to mediate in part the DM phenotype (6). Other substrates for DMPK include myogenin, L-type calcium channels, and Phospholemman (PLM) (1).

Pathways: [Regulation of Muscle Cell Differentiation](#), [Synaptic Membrane](#), [Skeletal Muscle Fiber Development](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months