antibodies -online.com





Datasheet for ABIN1393147

anti-IGHMBP2 antibody (AA 271-355) (Biotin)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	IGHMBP2	
Binding Specificity:	AA 271-355	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IGHMBP2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SMUBP2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig
Purification:	Purified by Protein A.

Target Details

Target:	IGHMBP2	
Alternative Name:	SMUBP2 (IGHMBP2 Products)	

Target Details

Background:

Synonyms: AEP, Antreeze enhancer binding protein, ATP-dependent helicase IGHMBP2, Cardiac transcription factor 1, Cardiac transcription factor 1, CATF 1, CATF 1, DNA-binding protein SMUBP-2, GF-1, Glial factor 1, HCSA, HMN 6, HMN6, IGHMBP 2, Ighmbp2, Immunoglobulin mu binding protein 2, Immunoglobulin mu binding protein 2, Immunoglobulin mu-binding protein 2, Immunoglobulin S mu binding protein 2, Immunoglobulin S mu binding protein 2, RIPE3 b1, RIPE3b 1, RIPE3b1, SMARD 1, SMARD1, SMBP2_HUMAN, SMUBP 2. Background: IGHMBP2 is a 993 amino acid nuclear and cytoplasmic protein that is ubiquitously

expressed. Belonging to the DNA2/NAM7 helicase family, IGHMBP2 is a 5' to 3' helicase that unwinds RNA and DNA duplexes in an ATP-dependent reaction. IGHMBP2 also acts as a transcriptional regulator and is necessary for transcriptional activation of the flounder liver-type antifreeze protein gene. IGHMBP2 exists as a homooligomer and is part of the cytosolic ribonucleoprotein complex. Mutations in the gene encoding IGHMBP2 are suggested to lead to distal hereditary motor neuronopathy type 6 (HMN6), also known as spinal muscular atrophy distal autosomal recessive 1 (DSMA1) or spinal muscular atrophy with respiratory distress 1 (SMARD1). HMN6 is characterized by weakness and wasting of distal muscles of the anterior tibial and peroneal compartments of the legs and severe respiratory distress.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	

Handling

Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months