antibodies -online.com





anti-ATL1 antibody (AA 201-300) (Alexa Fluor 647)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	ATL1	
Binding Specificity:	AA 201-300	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ATL1 antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SPG3A/Atlastin	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	ATL1
Alternative Name:	SPG3A/Atlastin (ATL1 Products)

Target Details

Background:

Synonyms: AD FSP, atl1, ATLA1_HUMAN, Atlastin GTPase 1, Atlastin-1, Atlastin1, Brain specic GTP binding protein, Brain-specic GTP-binding protein, FSP1, GBP-3, GBP3, GTP-binding protein 3, Guanine nucleotide-binding protein 3, Guanylate binding protein 3, hGBP3, HSN1D, Spastic paraplegia 3 protein A, SPG 3A, SPG3, SPG3A.

Background: Atlastins are Golgi-localized, integral membrane proteins that function as

Background: Atlastins are Golgi-localized, integral membrane proteins that function as GTPases. The Atlastin proteins, also designated SPG3A and guanylate-binding protein 3, comprise a Dynamin superfamily that plays a role in axonal maintenance. Hereditary spastic paraplegia (HSP) is an inherited neurodegenerative disorder that is characterized by retrograde axonal degeneration. HSP primarily affects long corticospinal neurons and causes spastic lower extremity weakness. Spastin, a microtubule (MT)-severing AAA ATPase, is a binding partner of Atlastin that is involved in membrane dynamics. This Spastin/Atlastin binding may be involved in the biochemical pathway that leads to HSP development. Mutations in the Atlastin gene (SPG3A) account for approximately 10 % of all autosomal dominant HSPs, while mutations in the Spastin gene (SPG4) account for almost 40 %.

Application Details

Application Notes:	IF(IHC-P) 1	

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: Fo

For Research Use only

:50-200

Handling

i landling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	